

CORRECTED VERSION

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
24 July 2003 (24.07.2003)

PCT

(10) International Publication Number  
**WO 03/059916 A3**

(51) International Patent Classification<sup>7</sup>: C07F 5/02

(21) International Application Number: PCT/US03/00768

(22) International Filing Date: 9 January 2003 (09.01.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/347,811 10 January 2002 (10.01.2002) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:  
US 60/347,811 (CON)  
Filed on 10 January 2002 (10.01.2002)

(71) Applicant (for all designated States except US): THE PENNSYLVANIA STATE RESEARCH FOUNDATION [US/US]; 304 Old Main, University Park, PA 16802-1504 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): TAVASSOLI, Ali [GB/US]; 710 South Atherton, #410, State College, PA 16801 (US). BENKOVIC, Stephen, J. [US/US]; 771 Teaberry Lane, State College, PA 16803 (US).

(74) Agent: SHALTOUT, Raafat, M.; McDonnell Boehnen Hulbert Berghoff, 300 South Wacker Drive, Suite 3200, Chicago, IL 60606 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(48) Date of publication of this corrected version:

30 October 2003

(15) Information about Correction:

see PCT Gazette No. 44/2003 of 30 October 2003, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS FOR THE PREPARATION OF ALKYL DIARYL BORINATES AND COMPLEXED DIARYLBORINIC ACIDS

(57) Abstract: The present invention provides methods for preparing complexed diaryl borinic acids from alkyl diarylborinates.

WO 03/059916 A3

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
24 July 2003 (24.07.2003)

PCT

(10) International Publication Number  
**WO 2003/059916 A3**

(51) International Patent Classification<sup>7</sup>: **C07F 5/02**

(21) International Application Number:  
PCT/US2003/000768

(22) International Filing Date: 9 January 2003 (09.01.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/347,811 10 January 2002 (10.01.2002) US

(63) Related by continuation (CON) or continuation-in-part  
(CIP) to earlier application:  
US 60/347,811 (CON)  
Filed on 10 January 2002 (10.01.2002)

(71) Applicant (for all designated States except US): **THE  
PENNSYLVANIA STATE RESEARCH FOUNDA-  
TION** [US/US]; 304 Old Main, University Park, PA  
16802-1504 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **TAVASSOLI, Ali**  
[GB/US]; 710 South Atherton, #410, State College, PA  
16801 (US). **BENKOVIC, Stephen, J.** [US/US]; 771  
Teaberry Lane, State College, PA 16803 (US).

(74) Agent: **SHALTOUT, Raafat, M.**; McDonnell Boehnen  
Hulbert Berghoff, 300 South Wacker Drive, Suite 3200,  
Chicago, IL 60606 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN,  
YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI,  
SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,  
GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:  
8 January 2004

(15) Information about Correction:

Previous Correction:

see PCT Gazette No. 44/2003 of 30 October 2003, Sec-  
tion II

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: METHODS FOR THE PREPARATION OF ALKYL DIARYL BORINATES AND COMPLEXED DIARYLBORINIC  
ACIDS

(57) Abstract: The present invention provides methods for preparing complexed diaryl borinic acids from alkyl diarylborinates.

WO 2003/059916 A3

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 C07F5/02		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07F		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) BEILSTEIN Data, EPO-Internal, WPI Data, CHEM ABS Data		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ZHANG, GUOMIN ET AL: "Studies on antitumor boron compounds. III. Synthesis of diarylborane chelates containing an intramolecular boron-nitrogen coordination bond and their antitumor activity" YOUJI HUAXUE (1982), (6), 409-12 , XP009010099 abstract page 410, paragraph 2 <div style="text-align: center; margin-top: 20px;">-/--</div>	1-3,6,15
<div style="display: flex; justify-content: space-between;"> <span><input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.</span> <span><input checked="" type="checkbox"/> Patent family members are listed in annex.</span> </div>		
* Special categories of cited documents :		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*&amp;* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold; margin-top: 10px;">30 April 2003</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold; margin-top: 10px;">19/05/2003</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold; margin-top: 10px;">Bork, A-M</div>

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE CROSSFIRE BEILSTEIN 'Online! Beilstein Institut zur Förderung der Chemischen Wissenschaften, Frankfurt am Main, DE; Database accession no. 759640 (reaction ID) XP002239683 abstract & J.AMER.CHEM.SOC., vol. 80, 1958, page 4291	9,12,13, 16
X	DATABASE CROSSFIRE BEILSTEIN 'Online! Beilstein Institut zur Förderung der Chemischen Wissenschaften, Frankfurt am Main, DE; Database accession no. 762485 (reaction ID) XP002239684 abstract & IZV.AKAD.NAUK SSSR SER.KHIM, 1956, pages 451-453,	9,11-13, 16
X	DATABASE CROSSFIRE BEILSTEIN 'Online! Beilstein Institut zur Förderung der Chemischen Wissenschaften, Frankfurt am Main, DE; Database accession no. 983118 (reaction ID) XP002239685 abstract & J.ORGANOMET. CHEM., vol. 156, 1978, pages 101-110,	9,11-13, 16
X	WO 00 75142 A (WALL MARK ;BAKER STEPHEN J (US); SHAPIRO LUCILLE (US); WAHNON DAPH) 14 December 2000 (2000-12-14) cited in the application page 17, line 15-21 page 38, line 11 -page 40, line 5 page 43, line 1-5 claims 9,14	15,17
A	page 44, line 1-17	1
P,X	WO 02 44184 A (WAHNON DAPHNE C ;BAKER STEPHEN J (GB); WALL MARK (US); BABOVAL JUS) 6 June 2002 (2002-06-06) page 6, line 27 -page 7, line 17 claim 2	15,17
A	page 16, line 15 -page 18, line 5	1

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 0075142	A	14-12-2000	AU	7569800 A	28-12-2000
			CA	2373279 A1	14-12-2000
			CN	1370170 T	18-09-2002
			EP	1181291 A2	27-02-2002
			JP	2003501431 T	14-01-2003
			WO	0075142 A2	14-12-2000
<hr/>					
WO 0244184	A	06-06-2002	AU	3940702 A	11-06-2002
			WO	0244184 A2	06-06-2002
<hr/>					

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
24 July 2003 (24.07.2003)

PCT

(10) International Publication Number  
**WO 03/059916 A2**

- (51) International Patent Classification<sup>7</sup>: **C07F 5/00**
- (21) International Application Number: PCT/US03/00768
- (22) International Filing Date: 9 January 2003 (09.01.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/347,811 10 January 2002 (10.01.2002) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:  
US 60/347,811 (CON)  
Filed on 9 January 2002 (09.01.2002)
- (71) Applicant (*for all designated States except US*): **THE PENNSYLVANIA STATE RESEARCH FOUNDATION** [US/US]; 304 Old Main, University Park, PA 16802-1504 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **TAVASSOLI, Ali** [GB/US]; 710 South Atherton, #410, State College, PA 16801 (US). **BENKOVIC, Stephen, J.** [US/US]; 771 Teaberry Lane, State College, PA 16803 (US).
- (74) Agent: **SHALTOUT, Raafat, M.**; McDonnell Boehnen Hulbert Berghoff, 300 South Wacker Drive, Suite 3200, Chicago, IL 60606 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: METHODS FOR THE PREPARATION OF ALKYL DIARYL BORINATES AND COMPLEXED DIARYLBORONIC ACIDS

(57) Abstract: The present invention provides methods for preparing complexed diaryl boronic acids from alkyl diarylborinates.

WO 03/059916 A2

# **METHODS FOR THE PREPARATION OF ALKYL DIARYL BORINATES AND COMPLEXED DIARYLBORONIC ACIDS**

## **CROSS-REFERENCE TO RELATED APPLICATIONS**

This application claims priority from U.S. Provisional Application Serial Number 60/347,811, filed January 9, 2002, the disclosure of which is incorporated herein by reference in its entirety.

## **BACKGROUND OF THE INVENTION**

### **1. Field of the Invention**

This invention relates to methods for preparing alkyl diarylborinates and methods for preparing complexed diarylboronic acids, which are useful as antibacterial agents.

### **2. Description of the Related Art**

N-O complexed diarylboronic acids are known to be effective pharmaceutical agents. For example, such compounds are capable of inhibiting adenine DNA methyltransferases in bacterial cells and thus exhibit antibacterial, growth-inhibitory properties against any bacterial species that produces an adenine DNA methyltransferase. Antibacterial diarylboronic acids are described in International Application Publication No. WO 00/75142.


Currently available methods for the preparation of N-O complexed diarylboronic acids require the use of dichloroborane methyl sulfide complex as a starting material. The use of dichloroborane methyl sulfide complex has several disadvantages, including the formation of dimethyl sulfide (DMS) as a side product of the reaction. Further, dichloroborane methyl sulfide reacts violently with water to produce flammable and harmful gases and has a repugnant odor that causes great discomfort for any lab personnel working with the material.

The formation of DMS as a side product makes the process of isolating a reaction product without contaminating the working area with DMS extremely difficult. Like dichloroborane methyl sulfide complex, DMS also has a foul odor. Further, DMS is a cancer suspect agent.

A need exists, therefore, for a method for preparing N-O complexed diarylboronic acids, and complexed diarylboronic acids in general, that is safer, more versatile and friendlier to lab personnel than currently known methods.


## SUMMARY OF THE INVENTION

The present invention provides a method for preparing a compound of the formula I:



The invention also provides a method for preparing a compound of formula

III:




The invention further provides compounds of formula I and formula III prepared according to the methods of the invention.

The methods of the invention are advantageous, *inter alia*, because they permit efficient synthesis of alkyl diarylborinates using reagents that are less toxic, less noxious and less dangerous than conventional reagents used in these syntheses.

Specific preferred embodiments of the present invention will become evident from the following more detailed description of certain preferred embodiments and the claims.

### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

As indicated above, the invention provides methods of preparing compounds of formula I:



wherein A is N, O or S;

W is C<sub>p</sub>, where p is 0 or 1;

R<sup>a</sup>, R<sup>b</sup>, R<sup>c</sup>, R<sup>d</sup>, and R<sup>e</sup> are the same or different and are independently hydrogen, halogen, nitro, nitroso, lower alkyl, aryl or substituted aryl, lower alkoxy, lower alkoxyalkyl, or cycloalkyl or cycloalkyl alkoxy, where each cycloalkyl group has from 3-7 members, where up to two of the cycloalkyl members are optionally hetero atoms selected from sulfur, oxygen and nitrogen, and where any member of the alkyl, aryl

or cycloalkyl group is optionally substituted with halogen, lower alkyl or lower alkoxy, aryl or substituted aryl, halogen, nitro, nitroso, aldehyde, carboxylic acid, amide, ester, or sulfate, or wherein  $R^a$ ,  $R^b$ ,  $R^c$ ,  $R^d$ , and  $R^e$  may be connected by aromatic, aliphatic, heteroaromatic, heteroaliphatic ring structures or substituted embodiments thereof, where  $R^a$  is absent when A is O or S and  $R^d$  is absent when  $p = 0$ ;

$R^f$  is hydrogen or is absent; and

wherein  $Ar^1$  and  $Ar^2$  can be the same or different and are each independently thienyl, aryl or aryl substituted at one or a plurality of positions with halogen, nitro, nitroso, lower alkyl, aryl or substituted aryl, lower alkoxy, lower alkoxyalkyl, or cycloalkyl or cycloalkyl alkoxy, where each cycloalkyl group has from 3-7 members, where up to two of the cycloalkyl members are optionally hetero atoms selected from sulfur, oxygen and nitrogen, and where any member of the alkyl, aryl or cycloalkyl group is optionally substituted with halogen, lower alkyl or lower alkoxy, aryl or substituted aryl, halogen, nitro, nitroso, aldehyde, carboxylic acid, amide, ester, or sulfate, and



wherein bond 1, bond 2, bond 3 and bond 4 are independently a single bond or a double bond, provided that when A is S or O, bond 1 is a single bond and where A is N, bond 1 is a double bond.

Compounds of formula I are useful as pharmaceutical agents. For example, compounds of formula I are antibacterial compounds, as described in International Application Publication No. WO 00/75142, and pending U.S. patent application Serial

No. 09/578,991, filed May 25, 2000, each of which is incorporated herein by reference in its entirety.




The methods of the invention for preparing compounds of formula I are depicted in Reaction Schemes 1 and 2.

### REACTION SCHEME 1



M is, for example, MgBr or Li

### REACTION SCHEME 2



Referring to Reaction Scheme 1, a trialkylborate is reacted with metalloorganic reagents  $\text{Ar}^1\text{M}$  and  $\text{Ar}^2\text{M}$  to form an alkyl diarylborinate of formula III. Preferably, the reaction is conducted by the addition of the trialkylborate to the metalloorganic reagents in an aprotic, non-aqueous solvent, such as tetrahydrofuran (THF) or diethyl ether. Also preferably, the addition is conducted under an inert atmosphere, such as a nitrogen or noble gas atmosphere, and at a temperature lower than room temperature, more preferably a temperature lower than 0 °C. For example, the addition is advantageously conducted at -78 °C under argon gas.

Following completion, the reaction is worked up by either a non-acidic workup step or by an acidic workup step. A non-acidic workup is preferred. In a non-acidic workup, the reaction mixture is treated with an excess of an organic solvent, most preferably an alcohol solvent such as methanol. The product is then isolated from the reaction mixture by known methods, such as by removal of the solvent under vacuum, and then the alkyl boronate is extracted with an ether or other organic solvent from the aqueous metal salts from the metalloorganic reagent. In the less preferred acidic workup step, an acidic solvent, such as dilute hydrochloric acid, is added to the reaction mixture to give a diaryl boronic acid ( $\text{Ar}^1\text{Ar}^2\text{B-OH}$ ). The boronic acid is then converted to the borinate of formula III by treating the boronic acid with an alcohol such as methanol.


Metalloorganic reagents suitable for use in the methods of the invention include Grignard reagents and organic alkali metal compounds, wherein the alkali metal is lithium or sodium. Other metalloorganic reagents are known in the art, such as zinc, copper and lithium metalloorganic reagents, can be used. A preferred metalloorganic reagent is a Grignard reagent. Grignard reagents are well known, and many are commercially available. Others can be readily prepared by known methods.

At least two equivalents of metalloorganic reagent are used for each equivalent of alkylborate. A single metalloorganic reagent can be used (*i.e.*, where  $\text{Ar}^1\text{M}$  and  $\text{Ar}^2\text{M}$  in Scheme 1 are the same), thus yielding an alkyl diarylborinate  $\text{Ar}^1\text{Ar}^2\text{B-O-alkyl}$  in which  $\text{Ar}^1$  and  $\text{Ar}^2$  are the same. Alternatively, two different metalloorganic reagents can be used, yielding an alkyl diarylborinate in which  $\text{Ar}^1$  and  $\text{Ar}^2$  are not the same.


Any alkylborate,  $\text{B(O-Alkyl)}_3$ , is suitable for use in the reaction described above. Preferred alkylborates include lower alkyl boronates, comprising 1-6 carbon atoms per alkyl group. Examples of preferred alkylboronates include but are not limited to trimethylborate, triethylborate, tributylborate or mixtures thereof. A more preferred alkylborate is trimethylborate. Alkylborates are commercially available (for example from Sigma-Aldrich, Milwaukee, Wisconsin) or can be readily prepared by known methods.

Referring to Reaction Scheme 2, the alkyl diarylborinate of formula III prepared as described in Reaction Scheme 1 is complexed with a complexing agent of formula II. Preferably, the complexing step is performed by dissolving or dispersing the complexing agent of formula II in a solvent and adding the solution or dispersion to a solution or dispersion of the alkyl diarylborinate of formula III. The product compound of formula I can be isolated by a variety of techniques including crystallization and filtration, or removal of solvent in vacuum. The product can be purified by known methods, including recrystallization, and/or chromatography.


The complexing agent II can be any compound containing an amino group and either a hydroxy, a thiol or a second amino group. The amino group and the hydroxy, thiol or second amino group are separated by 2 or 3 carbon atoms. Examples of such complexing agents include:



optionally substituted 8-hydroxyquinolines



optionally substituted *o*-amino phenols



amino alcohols or amino acids

wherein,

*n* is 1 or 2;

R and R' are the same or different and are independently hydrogen, halogen, lower

alkyl or lower alkoxy, aryl or substituted aryl, halogen, nitro, nitroso,

aldehyde, carboxylic acid, amide, ester, or sulfate, and

R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are the same or different and are independently hydrogen, halogen,

nitro, nitroso, lower alkyl, aryl or substituted aryl, lower alkoxy, lower

alkoxyalkyl, or cycloalkyl or cycloalkyl alkoxy, where each cycloalkyl group

has from 3-7 members, where up to two of the cycloalkyl members are

optionally hetero atoms selected from sulfur, oxygen and nitrogen, and where

any member of the alkyl, aryl or cycloalkyl group is optionally substituted

with halogen, lower alkyl or lower alkoxy, aryl or substituted aryl, halogen,


nitro, nitroso, aldehyde, carboxylic acid, amide, ester, or sulfate, or R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>,

and  $R_4$  may be connected by aromatic, aliphatic, heteroaromatic, heteroaliphatic ring structures or substituted embodiments thereof.


Preferred compounds of Formula I prepared according to the methods of the invention include those wherein A is oxygen.

Preferred compounds of Formula I prepared according to the methods of the invention also include those wherein  $Ar^1 = Ar^2$ .

Preferred compounds of Formula I prepared according to the methods of the invention also include those wherein  $Ar^1$  and  $Ar^2$  are selected from:



Preferred compounds of Formula II prepared according to the methods of the invention include:



The methods of the invention have several distinct advantages over other procedures for the preparation of complexed diarylboronic acids. For example, by using a trialkylborate as a starting material, chloroborane methyl sulfide is not required and the concomitant formation of DMS as a side product is avoided. As a

result, the process of isolating a reaction product is greatly simplified. Additionally, when trialkylborate reacts with water it produces alcohols, which are relatively harmless substances compared with the reaction side products produced according to the prior art.

In an industrial setting where large quantities of reagents are required, trialkylborinates are more appealing starting materials than the previously used reagents, not only because of the inherent safety of the borinates, but also because of their cost. Trimethylborate, for example, is considerably less expensive than previously used reagents.

The presence of an alkyl ester in compounds of formula III allows this compound to form a complex with a wider variety of complexing agents of formula II than was previously possible. This is due in part to increased solubility of the compound of formula III and because there is a decreased possibility of side reactions and side products using these compounds. Previous methods for synthesizing complexed diarylboronic acids used a boronic acid in the complexing step. Boronic acids, however, can react with any amine (base) in the complexing agent to produce a salt, which produces an undesired product. The boronic esters of formula III of the invention, on the other hand, can react with a wider variety of compounds, including compounds containing hydrophilic or polar charged substituents, without producing such undesired products.

A further advantage of the methods of the invention is that diarylborinates of formula III in the invention are prepared in a single step. Prior art methods prepared boronic acid in a first step and required a second step to produce the alkyl boronate, and are thus less robust than the instant methods.

Compounds of Formula I prepared according to the methods of the invention may contain one or more asymmetric carbon atoms, so that the compounds can exist in different stereoisomeric forms. These compounds can be, for example, racemates or optically active forms. In these situations, the single enantiomers, *i.e.*, optically active forms, can be obtained by asymmetric synthesis or by resolution of the racemates. Resolution of the racemates can be accomplished, for example, by conventional methods such as crystallization in the presence of a resolving agent, or chromatography using, for example a chiral HPLC column.

Compounds prepared according to the methods of the invention can exist as tautomers in solution. When structures and names are given for one tautomeric form the other tautomeric form is also included in the invention.

Representative compounds prepared according to the methods of the present invention include, but are not limited to the compounds disclosed herein and their pharmaceutically acceptable acid and base addition salts. In addition, if a compound is obtained as an acid addition salt, the free base can be obtained by basifying a solution of the acid salt. Conversely, if the product is a free base, an addition salt, particularly a pharmaceutically acceptable addition salt, may be produced by dissolving the free base in a suitable organic solvent and treating the solution with an acid, in accordance with conventional procedures for preparing acid addition salts from base compounds.

In the description of the reagents and methods of this invention, particular terms are defined as follows.

By "alkyl", "lower alkyl", and "C<sub>1</sub>-C<sub>6</sub> alkyl" in the present invention is meant straight or branched chain alkyl groups having 1-6 carbon atoms, such as, methyl,

ethyl, propyl, isopropyl, *n*-butyl, *sec*-butyl, *tert*-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, and 3-methylpentyl.

By "alkoxy", "lower alkoxy", and "C<sub>1</sub>-C<sub>6</sub> alkoxy" in the present invention is meant straight or branched chain alkoxy groups having 1-6 carbon atoms, such as, for example, methoxy, ethoxy, propoxy, isopropoxy, *n*-butoxy, *sec*-butoxy, *tert*-butoxy, pentoxy, 2-pentyl, isopentoxy, neopentoxy, hexoxy, 2-hexoxy, 3-hexoxy, and 3-methylpentoxy.

By the term "halogen" in the present invention is meant fluorine, bromine, chlorine, and iodine.

By "aliphatic ring" or "cycloalkyl", e.g., C<sub>3</sub>-C<sub>7</sub> cycloalkyl, in the present invention is meant cycloalkyl groups having 3-7 atoms such as, for example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl. In the C<sub>3</sub>-C<sub>7</sub> cycloalkyl groups, preferably in the C<sub>5</sub>-C<sub>7</sub> cycloalkyl groups, one or two of the carbon atoms forming the ring can optionally be replaced with a heteroatom, such as sulfur, oxygen or nitrogen. Examples of such groups are piperidiny, piperaziny, morpholiny, pyrrolidiny, imidazolidiny, oxazolidiny, azaperhydroepiny, oxazaperhydroepiny, oxepany, oxazaperhydroiny, and oxadiazaperhydroiny. C<sub>3</sub> and C<sub>4</sub> cycloalkyl groups having a member replaced by nitrogen or oxygen include aziridiny, azetidiny, oxetany, and oxirany.

By "aryl" is meant an aromatic carbocyclic group having a single ring (e.g., phenyl), multiple rings (e.g., biphenyl), or multiple condensed rings in which at least one is aromatic, (e.g., 1,2,3,4-tetrahydronaphthyl, naphthyl, anthryl, or phenanthryl), which is optionally mono-, di-, or trisubstituted with, e.g., halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, lower acyloxy, aryl, heteroaryl, and hydroxy.

Preferred aryl groups include phenyl and naphthyl, each of which is optionally substituted as defined herein.

By "heteroaromatic" or "heteroaryl" is meant one or more aromatic ring systems of 5-, 6-, or 7-membered rings containing at least one and up to four heteroatoms selected from nitrogen, oxygen, or sulfur. Such heteroaryl groups include, for example, thienyl, furanyl, thiazolyl, imidazolyl, (is)oxazolyl, pyridyl, pyrimidinyl, (iso)quinolinyl, naphthyridinyl, benzimidazolyl, and benzoxazolyl. Preferred heteroaryls are thiazolyl, pyrimidinyl, preferably pyrimidin-2-yl, and pyridyl. Other preferred heteroaryl groups include 1-imidazolyl, 2-thienyl, 1-, or 2-quinolinyl, 1-, or 2-isoquinolinyl, 1-, or 2-tetrahydroisoquinolinyl, 2- or 3-furanyl and 2-tetrahydrofuranyl.

By "aliphatic ring," or "heterocycle," is meant one or more carbocyclic ring systems of 3-, 4-, 5-, 6-, or 7-membered rings which includes fused ring systems of 9-11 atoms containing at least one and up to four heteroatoms selected from nitrogen, oxygen, or sulfur. Preferred heterocycles of the present invention include morpholinyl, thiomorpholinyl, thiomorpholinyl S-oxide, thiomorpholinyl S,S-dioxide, piperazinyl, homopiperazinyl, pyrrolidinyl, pyrrolinyl, tetrahydropyranyl, piperidinyl, tetrahydrofuranyl, tetrahydrothienyl, homopiperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S,S-dioxide, oxazolidinonyl, dihydropyrazolyl, dihydropyrrolyl, dihydropyrazinyl, dihydropyridinyl, dihydropyrimidinyl, dihydrofuryl, dihydropyranyl, azepanyl, diazepanyl, tetrahydrothienyl S-oxide, tetrahydrothienyl S,S-dioxide and homothiomorpholinyl S-oxide.

Compounds prepared according to the methods of the invention can be provided as pharmaceutical compositions. The pharmaceutical compositions can be

manufactured in a manner that is itself known, *e.g.*, by means of a conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Pharmaceutical compositions can be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries that facilitate processing of the active compounds into preparations that can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

Non-toxic pharmaceutical salts include salts of acids such as hydrochloric, phosphoric, hydrobromic, sulfuric, sulfinic, formic, toluenesulfonic, methanesulfonic, nitic, benzoic, citric, tartaric, maleic, hydroiodic, alkanoic such as acetic,  $\text{HOOC}-(\text{CH}_2)_n-\text{CH}_3$  where  $n$  is 0-4, and the like. Non-toxic pharmaceutical base addition salts include salts of bases such as sodium, potassium, calcium, ammonium, and the like. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable addition salts.

For injection, the compounds prepared according to the methods of the invention can be formulated in appropriate aqueous solutions, such as physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal and transcutaneous administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries,

suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents can be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions can be used, which can optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments can be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations that can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds can be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers can be added. All formulations for oral administration should be in dosages suitable for such

administration. For buccal administration, the compositions can take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds prepared according to the methods of the invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of *e.g.*, gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds can be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection can be presented in unit dosage form, *e.g.*, in ampoules or in multi-dose containers, with an added preservative. The compositions can take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds can be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions can contain substances that increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension can also contain suitable stabilizers or agents that increase

the solubility of the compounds to allow for the preparation of highly concentrated solutions. Alternatively, the active ingredient can be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use. The compounds can also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds can also be formulated as a depot preparation. Such long acting formulations can be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

A pharmaceutical carrier for hydrophobic compounds of formula I is a cosolvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. The cosolvent system can be the VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant polysorbate 80, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:5W) consists of VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system can be varied considerably without destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components can be varied: for example, other low-toxicity nonpolar surfactants can be used instead of polysorbate 80; the

fraction size of polyethylene glycol can be varied; other biocompatible polymers can replace polyethylene glycol, *e.g.* polyvinyl pyrrolidone; and other sugars or polysaccharides can substitute for dextrose.

Alternatively, other delivery systems for hydrophobic pharmaceutical compounds can be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also can be employed, although usually at the cost of greater toxicity. Additionally, the compounds can be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules can, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein and nucleic acid stabilization can be employed.

The pharmaceutical compositions also can comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

The compounds of Formula I can be provided as salts with pharmaceutically compatible counterions. Pharmaceutically compatible salts can be formed with many acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, phosphoric, hydrobromic, sulfinic, formic, toluenesulfonic, methanesulfonic, nitric, benzoic, citric, tartaric, maleic, hydroiodic, alkanoic such as acetic,  $\text{HOOC}-(\text{CH}_2)_n-\text{CH}_3$  where  $n$  is 0-4, and the like. Salts tend to be more soluble in aqueous or

other protonic solvents that are the corresponding free base forms. Non-toxic pharmaceutical base addition salts include salts of bases such as sodium, potassium, calcium, ammonium, and the like. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable addition salts.

Pharmaceutical compositions of the compounds prepared according to the methods of the invention can be formulated and administered through a variety of means, including systemic, localized, or topical administration. Techniques for formulation and administration can be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA. The mode of administration can be selected to maximize delivery to a desired target site in the body. Suitable routes of administration can, for example, include oral, rectal, transmucosal, transcutaneous, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.

Alternatively, one can administer the compound in a local rather than systemic manner, for example, *via* injection of the compound directly into a specific tissue, often in a depot or sustained release formulation.

Pharmaceutical compositions suitable for use include compositions wherein the active ingredients are contained in an effective amount to achieve its intended purpose. More specifically, a therapeutically effective amount means an amount effective to prevent development of or to alleviate the existing symptoms of the subject being treated. Determination of the effective amounts is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

For administration to non-human animals, the drug or a pharmaceutical composition containing the drug may also be added to the animal feed or drinking water. It will be convenient to formulate animal feed and drinking water products with a predetermined dose of the drug so that the animal takes in an appropriate quantity of the drug along with its diet. It will also be convenient to add a premix containing the drug to the feed or drinking water approximately immediately prior to consumption by the animal.

Preferred compounds prepared according to the methods of the invention will have certain pharmacological properties. Such properties include, but are not limited to oral bioavailability, low toxicity, low serum protein binding and desirable *in vitro* and *in vivo* half-lives. Assays may be used to predict these desirable pharmacological properties. Assays used to predict bioavailability include transport across human intestinal cell monolayers, including Caco-2 cell monolayers. Serum protein binding may be predicted from albumin binding assays. Such assays are described in a review by Oravcová *et al.* (1996, *J. Chromat. B* 677: 1-27). Compound half-life is inversely proportional to the frequency of dosage of a compound. *In vitro* half-lives of compounds may be predicted from assays of microsomal half-life as described by Kuhnz and Gieschen (1998, DRUG METABOLISM AND DISPOSITION, vol. 26, pp. 1120-1127).

Toxicity and therapeutic efficacy of such compounds can be determined by conventional pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD50 and ED50. Compounds that exhibit high therapeutic indices are


preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED<sub>50</sub> with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See, e.g. Fingl *et al.*, 1975, in THE PHARMACOLOGICAL BASIS OF THERAPEUTICS, Ch.1, p.1).

Dosage amount and interval can be adjusted individually to provide plasma levels of the active moiety that are sufficient to maintain bacterial cell growth-inhibitory effects. Usual patient dosages for systemic administration range from 100 - 2000 mg/day. Stated in terms of patient body surface areas, usual dosages range from 50 - 910 mg/m<sup>2</sup>/day. Usual average plasma levels should be maintained within 0.1-1000 :M. In cases of local administration or selective uptake, the effective local concentration of the compound cannot be related to plasma concentration.


The disclosures in this application of all articles and references, including patents, are incorporated herein by reference.

The following Examples are provided for the purposes of illustration and are not intended to limit the scope of the present invention. The present invention is not to be limited in scope by the exemplified embodiments, which are intended as illustrations of individual aspects of the invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

## EXAMPLES


Example 1: General procedure for the synthesis of methyl diarylborinates

Trimethylborate (0.95 eq) is added dropwise to a freshly prepared solution of aryl magnesium bromide (2 eq) in tetrahydrofuran (0.3 M) under argon at  $-78\text{ }^{\circ}\text{C}$ . The mixture is warmed to room temperature and stirred overnight. The solution is cooled to  $-78\text{ }^{\circ}\text{C}$  and excess Grignard reagent is destroyed by the dropwise addition of methanol until no more effervescence is observed. Solvents are removed *in vacuo* and the residue dissolved in diethyl ether followed by a water wash. The organic layer is dried (over  $MgSO_4$ ), filtered and the solvents removed *in vacuo* to give the product as a foam that can be used without further purification. Any trialkylborate may be used as a substitute for trimethylborate including triethylborate and tributylborate, as discussed above.

Example 2: General procedure for the synthesis of N-O complexed diarylboronic acids

The complexing agent (0.9 eq) is dissolved in methanol (or dichloromethane depending on solubility) and added to a solution of methyl diarylborinate (1 eq) (prepared as outlined above) in methanol. The solution is allowed to stand overnight, if the product has crystallized, the solid is collected and washed with cold ethanol; otherwise the solvents are removed *in vacuo* to give a solid which is purified by recrystallization from ethanol or diethyl ether (depending on the complexing agent). If this is unsuccessful the product is purified by column chromatography on silica gel eluting with diethyl ether/hexane.


Example 3: Synthesis of methyl di(*p*-chlorophenyl) borinate



Trimethylborate (2.2 ml,  $1.92 \times 10^{-2}$  mol) is added dropwise to a freshly prepared solution of *p*-chlorophenyl magnesium bromide (40.4 ml, 1 M,  $4.04 \times 10^{-2}$  mol) in tetrahydrofuran (60 ml) under argon at  $-78^\circ\text{C}$ . The mixture is warmed to room temperature and stirred overnight. The solution is cooled to  $-78^\circ\text{C}$  and excess grignard reagent is destroyed by the dropwise addition of methanol until no more effervescence is observed. The solvents are removed *in vacuo* and the residue dissolved in diethyl ether and washed with water. The organic layer is dried ( $\text{MgSO}_4$ ),


filtered and the solvents removed *in vacuo* to give methyl di(*p*-chlorophenyl)borinate (4.60 g, 92 %) as a solid which was used without further purification.

Example 4: Synthesis of methyl di(*p*-fluorophenyl) borinate




Trimethylborate (1.0 ml,  $8.75 \times 10^{-3}$  mol) is added dropwise to a freshly prepared solution of *p*-fluorophenyl magnesium bromide (18.4 ml, 1 M,  $1.84 \times 10^{-2}$  mol) in tetrahydrofuran (30 ml) under argon at  $-78^\circ\text{C}$ . The mixture is warmed to room temperature and stirred overnight. The solution is cooled to  $-78^\circ\text{C}$  and excess grignard reagent is destroyed by the dropwise addition of methanol until no more effervescence is observed. The solvents are removed *in vacuo* and the residue dissolved in diethyl ether and washed with water. The organic layer is dried ( $\text{MgSO}_4$ ), filtered and the solvents removed *in vacuo* to give methyl di(*p*-fluorophenyl)borinate (1.93 g, 89 %) as a solid which was used without further purification.

Example 5: Synthesis of methyl di(*m*-chlorophenyl) borinate




Trimethylborate (1.3 ml,  $1.14 \times 10^{-2}$  mol) is added dropwise to a freshly prepared solution of *m*-chlorophenyl magnesium bromide (48 ml, 0.5 M,  $2.39 \times 10^{-2}$  mol) in tetrahydrofuran (30 ml) under argon at  $-78^\circ\text{C}$ . The mixture is warmed to room temperature and stirred overnight. The solution is cooled to  $-78^\circ\text{C}$  and excess grignard reagent is destroyed by the dropwise addition of methanol until no more effervescence is observed. The solvents are removed *in vacuo* and the residue dissolved in diethyl ether and washed with water. The organic layer is dried ( $\text{MgSO}_4$ ), filtered and the solvents removed *in vacuo* to give di(*m*-chlorophenyl)borinate (3.2 g, 94 %) as a solid which was used without further purification.

Example 6: Di(*p*-chlorophenyl)borinic acid 5-nitro-8-hydroxyquinoline ester




5-Nitro-8-hydroxyquinoline (340 mg,  $1.79 \times 10^{-3}$ ) was added to a solution of methyl di(*p*-chlorophenyl)borinate (500 mg,  $1.89 \times 10^{-3}$ ) in dichloromethane. The solution turned dark yellow and was left to stir overnight. The solvent was removed *in vacuo* to give a solid, which was washed with diethyl ether (20 ml) and recrystallized from ethanol to give the title compound (630 mg, 83 %) as a dark yellow solid.


**Example 7:** Di-(*m*-chlorophenyl)borinic acid 8-hydroxyquinoline ester



A solution of 8-hydroxyquinoline (2.03 g,  $1.4 \times 10^{-2}$  mol) in ethanol (40 ml) was added dropwise to methyl di(*m*-chlorophenyl) borinate (3.7 g,  $1.4 \times 10^{-2}$  mol) in ethanol (40 ml) at room temperature. A yellow precipitation was observed and was left to stand overnight. The solid was collected by filtration and washed with cold ethanol to yield the title product as a yellow solid; mp 144 – 145 °C ;  $^1\text{H-NMR}$  (360 MHz,  $\text{C}^2\text{H}_3\text{O}^2\text{H}$ ):  $\delta$  8.83 (d,  $J = 5.0$  Hz, 1H), 8.63 (d,  $J = 8.2$  Hz, 1H), 7.78 (dd,  $J = 8.6, 5.0$  Hz, 1H), 7.67 (t,  $J = 8.2$ , 1H), 7.36 (d,  $J = 8.2$ , 1H), 7.26-7.09 (m, 9H); MS (+ve ESI)  $m/z$  377 ( $[\text{M}+\text{H}]^+$ ,  $^{10}\text{B}$ ,  $^{35}\text{Cl}$ ,  $^{35}\text{Cl}$ ), 378 ( $[\text{M}+\text{H}]^+$ ,  $^{11}\text{B}$ ,  $^{35}\text{Cl}$ ,  $^{35}\text{Cl}$ ), 379 ( $[\text{M}+\text{H}]^+$ ,  $^{10}\text{B}$ ,  $^{35}\text{Cl}$ ,  $^{37}\text{Cl}$ ), 380 ( $[\text{M}+\text{H}]^+$ ,  $^{11}\text{B}$ ,  $^{35}\text{Cl}$ ,  $^{37}\text{Cl}$ ), 381 ( $[\text{M}+\text{H}]^+$ ,  $^{10}\text{B}$ ,  $^{37}\text{Cl}$ ,  $^{37}\text{Cl}$ ), 382 ( $[\text{M}+\text{H}]^+$ ,  $^{11}\text{B}$ ,  $^{37}\text{Cl}$ ,  $^{37}\text{Cl}$ ); Anal. ( $\text{C}_{21}\text{H}_{14}\text{NOBCl}_2$ ) C, H, N.

Example 8: Di-(*p*-chlorophenyl)borinic acid glycine ester


A solution of glycine (67 mg,  $8.99 \times 10^{-4}$  mol) in water (5 ml) was added dropwise to a solution of methyl di(*p*-chlorophenyl) borinate (250 mg,  $9.47 \times 10^{-4}$  mol) in ethanol (5 ml). The mixture was left to stir overnight. The solvents were removed *in vacuo* to give a solid, which was washed with diethyl ether (40 ml) and recrystallized from ethanol to give the title compound (225 mg, 82 %) as a white solid.

Example 9: Di-(*p*-chlorophenyl)borinic acid (L)-proline ester

A solution of (L)-proline (54 mg,  $4.69 \times 10^{-4}$  mol) in water (5 ml) was added dropwise to a solution of methyl di(*p*-chlorophenyl) borinate (130 mg,  $4.92 \times 10^{-4}$  mol) in ethanol (5 ml). The mixture was left to stir overnight. The solvents were


removed *in vacuo* to give a solid, which was washed with diethyl ether (40 ml) and recrystallized from ethanol to give the title compound (149 mg, 90 %) as a white solid.

**Example 10:** Di-(*p*-chlorophenyl)borinic acid N-hydroxyethyl cytosine ester



A solution of N-hydroxyethyl cytosine (100 mg,  $7.04 \times 10^{-4}$  mol) in water (5 ml) was added dropwise to a solution of methyl di(*p*-chlorophenyl) borinate (205 mg,  $7.75 \times 10^{-4}$  mol) in ethanol (5 ml). The mixture was left to stir overnight. The solvents were removed *in vacuo* to give a solid, which was washed with diethyl ether (30 ml) and recrystallized from ethanol to give the title compound (211 mg, 79 %) as a white solid.

**Example 11:** Di-(*p*-chlorophenyl)borinic acid N-hydroxyethyl 5-fluorocytosine ester




A solution of N-hydroxyethyl-5-fluorocytosine (364 mg,  $2.27 \times 10^{-3}$  mol) in water (5 ml) was added dropwise to a solution of methyl di(*p*-chlorophenyl) borinate (659 mg,  $2.50 \times 10^{-3}$  mol) in ethanol (20 ml). The mixture was left to stir overnight. The solvents were removed *in vacuo* to give a solid, which was washed with diethyl ether (40 ml) and recrystallized from ethanol to give the title compound (773 mg, 86 %) as a white solid.

The following compounds are prepared according to the methods of the invention:

Example  
No.


Compound

12




di-(*p*-fluorophenyl)borinic acid 8-hydroxyquinoline ester

13




di-(*p*-chlorophenyl)borinic acid 8-hydroxyquinoline ester

14




diphenylborinic acid 8-hydroxyquinoline ester

15




di-(p-fluorophenyl)borinic acid ethanolamine ester

16




di-(p-chlorophenyl)borinic acid ethanolamine ester

17




6-N-(diphenylborinic ester)-ethyl-adenine

18




6-N-(diphenylborinic ester)-ethyl-9-(2-(4-morpholinyl)-ethyl)-adenine

19




6-N-(diphenylborinic ester)-ethyl-9-(3-(N-phthaloyl)-aminopropyl)-  
adenine

20




6-N-(diphenylborinic ester)-ethyl-9-(2-(2-(2-hydroxyethoxy)ethoxy)-  
ethyl)-adenine

21




6-N-(diphenylborinic ester)-ethyl-9-(ethyl-2-acrylate)-methyl-adenine

22




## Di-(4-chloro-2-fluorophenyl)borinic acid 8-hydroxyquinoline ester

23




## Di-(3,4-methylenedioxyphenyl)borinic acid 8-hydroxyquinoline ester

24




## Di-(4-methoxyphenyl)borinic acid 8-hydroxyquinoline ester

25




## Di-(2-thienyl)borinic acid 8-hydroxyquinoline ester

26




## Di-(p-fluorophenyl)borinic acid 8-hydroxyquinoline ester

27




Di-(p-chlorophenyl)borinic acid 8-hydroxyquinoline ester

28




Di-(4-methoxyphenyl)borinic acid 8-hydroxyquinoline ester

29




Di-(p-fluorophenyl)borinic acid 5-chloro-8-hydroxyquinoline ester

30




Di-(p-chlorophenyl)borinic acid 5-chloro-8-hydroxyquinoline ester

31




Di-(3,4-methylenedioxyphenyl)borinic acid 5-chloro-8-hydroxyquinoline ester

32




Di-(4-methoxyphenyl)borinic acid 5-chloro-8-hydroxyquinoline ester

33




Di-(3,4-methylenedioxyphenyl)borinic acid 8-hydroxy-5-nitroquinoline ester

34




Diphenylborinic acid 2-aminophenol

35




Diphenylborinic acid pyridine-2-methanol

36




Diphenylborinic acid 2-amino-1-phenylpropanol

37




Diphenylborinic acid (S)-(+)-pyrrolidine-2-methanol

38




Di-(4-fluorophenyl)borinic acid ethanolamine ester

39




Di-(4-chlorophenyl)borinic acid ethanolamine ester

40




6-N-(diphenylborinic ester)-ethyl-9-(2-hydroxymethyl-5-methyl-tetrahydro-furan-3,4-diol)-adenine

41



6-amino-4(2-diphenylborinic ester) ethylamino pyrimidine

42




4-amino-5(2-diphenylborinic ester ethyliminoester)imidazole.

It should be understood that the foregoing disclosure emphasizes certain specific embodiments of the invention and that all modifications or alternatives equivalent thereto are within the spirit and scope of the invention as set forth in the appended claims.

## What is claimed is:

1. A method for preparing a compound of formula I:



wherein A is N, O or S;

W is C<sub>p</sub>, where p is 0 or 1;

R<sup>a</sup>, R<sup>b</sup>, R<sup>c</sup>, R<sup>d</sup>, and R<sup>e</sup> are the same or different and are independently hydrogen, halogen, nitro, nitroso, lower alkyl, aryl or substituted aryl, lower alkoxy, lower alkoxyalkyl, or cycloalkyl or cycloalkyl alkoxy, where each cycloalkyl group has from 3-7 members, where up to two of the cycloalkyl members are optionally hetero atoms selected from sulfur, oxygen and nitrogen, and where any member of the alkyl, aryl or cycloalkyl group is optionally substituted with halogen, lower alkyl or lower alkoxy, aryl or substituted aryl, halogen, nitro, nitroso, aldehyde, carboxylic acid, amide, ester, or sulfate, or wherein R<sup>a</sup>, R<sup>b</sup>, R<sup>c</sup>, R<sup>d</sup>, and R<sup>e</sup> may be connected by aromatic, aliphatic, heteroaromatic, heteroaliphatic ring structures or substituted embodiments thereof, where R<sup>a</sup> is absent when A is O or S, and R<sup>d</sup> is absent when p = 0;

R<sup>f</sup> is hydrogen or is absent; and



wherein Ar<sup>1</sup> and Ar<sup>2</sup> can be the same or different and are each independently thienyl, aryl or aryl substituted at one or a plurality of positions with

halogen, nitro, nitroso, lower alkyl, aryl or substituted aryl, lower alkoxy, lower alkoxyalkyl, or cycloalkyl or cycloalkyl alkoxy, where each cycloalkyl group has from 3-7 members, where up to two of the cycloalkyl members are optionally hetero atoms selected from sulfur, oxygen and nitrogen, and where any member of the alkyl, aryl or cycloalkyl group is optionally substituted with halogen, lower alkyl or lower alkoxy, aryl or substituted aryl, halogen, nitro, nitroso, aldehyde, carboxylic acid, amide, ester, or sulfate, and




wherein bond 1, bond 2, bond 3 and bond 4 are independently a single bond or a double bond, provided that when A is S or O, bond 1 is a single bond and where A is N, bond 1 is a double bond,

said method comprising the step of:

reacting an alkyl diarylborinate of formula III with a compound of formula II to form the compound of formula I



2. The method of claim 1 wherein the alkyl diarylborinate of formula III and the compound of formula II are in a ratio of about 1 to about 0.9 equivalents respectively.
3. The method of claim 1 wherein the compound of formula III is:



wherein,

n is 1 or 2;


R and R' are the same or different and are independently hydrogen, halogen, lower alkyl or lower alkoxy, aryl or substituted aryl, halogen, nitro, nitroso, aldehyde, carboxylic acid, amide, ester, or sulfate, and

R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> are the same or different and are independently hydrogen, halogen, nitro, nitroso, lower alkyl, aryl or substituted aryl, lower alkoxy, lower alkoxyalkyl, or cycloalkyl or cycloalkyl alkoxy, where each cycloalkyl group has from 3-7 members, where up to two of the cycloalkyl members are optionally hetero atoms selected from sulfur, oxygen and nitrogen, and where any member of the alkyl, aryl or cycloalkyl group is optionally substituted with halogen, lower alkyl or lower alkoxy, aryl or substituted aryl, halogen, nitro, nitroso, aldehyde, carboxylic acid, amide, ester, or sulfate, or R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, and R<sub>4</sub> may be connected by aromatic, aliphatic, heteroaromatic, heteroaliphatic ring structures or substituted embodiments thereof.

4. The method of claim 1 wherein the alkyl diarylborinate of formula III is prepared by reacting a trialkylborate with a metalloorganic reagent.

5. The method of claim 4 wherein the trialkylborate is trimethylborate, triethylborate, tributylborate, or mixtures thereof.

6. The method of claim 4 wherein the metalloorganic reagent is a Grignard reagent or a lithium reagent.
7. The method of claim 4 wherein the trialkylborate and the metalloorganic reagent are in a ratio of about 1 to about 2 equivalents respectively.
8. The method of claim 4 further comprising the step of treating the reaction product with methanol.
9. A method for preparing a compound of formula III:



wherein  $\text{Ar}^1$  and  $\text{Ar}^2$  can be the same or different and are each independently aryl or aryl substituted at one or a plurality of positions with halogen, nitro, nitroso, lower alkyl, aryl or substituted aryl, lower alkoxy, lower alkoxyalkyl, or cycloalkyl or cycloalkyl alkoxy, where each cycloalkyl group has from 3-7 members, where up to two of the cycloalkyl members are optionally hetero atoms selected from sulfur, oxygen and nitrogen, and where any member of the alkyl, aryl or cycloalkyl group is optionally substituted with halogen, lower alkyl or lower alkoxy, aryl or substituted aryl, halogen, nitro, nitroso, aldehyde, carboxylic acid, amide, ester, or sulfate,

said method comprising the step of reacting a trialkylborate with a metalloorganic reagent.

11. The method of claim 9 wherein the trialkylborate is trimethylborate, triethylborate, tributylborate, or mixtures thereof.
12. The method of claim 9 wherein the metalloorganic reagent is a Grignard reagent or a lithium reagent.
13. The method of claim 9 wherein the trialkylborate and the metalloorganic reagent are in a ratio of about 1 to about 2 equivalents respectively.
14. The method of claim 9 further comprising the step of treating the reaction product with methanol.
15. A compound of the formula I prepared by the method of claim 1.
16. A compound of the formula III prepared by the method of claim 9.
17. A compound according to claim 15 that is:
  - Di-(*p*-chlorophenyl)borinic acid 5-nitro-8-hydroxyquinoline ester,
  - Di-(*m*-chlorophenyl)borinic acid 8-hydroxyquinoline ester,
  - Di-(*p*-chlorophenyl)borinic acid glycine ester,
  - Di-(*p*-chlorophenyl)borinic acid (L)-proline ester,
  - Di-(*p*-chlorophenyl)borinic acid N-hydroxyethyl cytosine ester,

Di-(*p*-chlorophenyl)borinic acid N-hydroxyethyl 5-fluorocytosine ester,  
di-(*p*-fluorophenyl)borinic acid 8-hydroxyquinoline ester,  
di-(*p*-chlorophenyl)borinic acid 8-hydroxyquinoline ester,  
diphenylborinic acid 8-hydroxyquinoline ester,  
di-(*p*-fluorophenyl)borinic acid ethanolamine ester,  
di-(*p*-chlorophenyl)borinic acid ethanolamine ester,  
6-N-(diphenylborinic ester)-ethyl-adenine,  
6-N-(diphenylborinic ester)-ethyl-9-(2-(4-morpholinyl)-ethyl)-adenine,  
6-N-(diphenylborinic ester)-ethyl-9-(3-(N-phthaloyl)-aminopropyl)-adenine,  
6-N-(diphenylborinic ester)-ethyl-9-(2-(2-(2-hydroxyethoxy)ethoxy)-ethyl)-  
adenine,  
6-N-(diphenylborinic ester)-ethyl-9-(ethyl-2-acrylate)-methyl-adenine,  
Di-(4-chloro-2-fluorophenyl)borinic acid 8-hydroxyquinoline ester,  
Di-(3,4-methylenedioxyphenyl)borinic acid 8-hydroxyquinoline ester,  
Di-(4-methoxyphenyl)borinic acid 8-hydroxyquinoline ester,  
Di-(2-thienyl)borinic acid 8-hydroxyquinoline ester,  
Di-(*p*-fluorophenyl)borinic acid 8-hydroxyquinaldine ester,  
Di-(*p*-chlorophenyl)borinic acid 8-hydroxyquinaldine ester,  
Di-(4-methoxyphenyl)borinic acid 8-hydroxyquinaldine ester,  
Di-(*p*-fluorophenyl)borinic acid 5-chloro-8-hydroxyquinoline ester,  
Di-(*p*-chlorophenyl)borinic acid 5-chloro-8-hydroxyquinoline ester,  
Di-(3,4-methylenedioxyphenyl)borinic acid 5-chloro-8-hydroxyquinoline  
ester,  
Di-(4-methoxyphenyl)borinic acid 5-chloro-8-hydroxyquinoline ester,  
Di-(3,4-methylenedioxyphenyl)borinic acid 8-hydroxy-5-nitroquinoline

ester,

Diphenylborinic acid 2-aminophenol,

Diphenylborinic acid pyridine-2-methanol,

Diphenylborinic acid 2-amino-1-phenylpropanol,

Diphenylborinic acid (S)-(+)-pyrrolidine-2-methanol,

Di-(4-fluorophenyl)borinic acid ethanolamine ester,

Di-(4-chlorophenyl)borinic acid ethanolamine ester.

6-N-(diphenylborinic ester)-ethyl-9-(2-hydroxymethyl-5-methyl-tetrahydro-furan-3,4-diol)-adenine,

6-amino-4(2-diphenylborinic ester) ethylamino pyrimidine, or

4-amino-5(2-diphenylborinic ester ethyliminoester)imidazole.



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>C07D 473/00, A61K 31/52,</b> <b>A61P 31/00, C07D 473/34,</b> <b>C07H 19/16, C07H 19/20</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/75142</b> <b>(43) International Publication Date:</b> 14 December 2000 (14.12.2000)
<b>(21) International Application Number:</b> PCT/US00/14479 <b>(22) International Filing Date:</b> 25 May 2000 (25.05.2000) <b>(30) Priority Data:</b> 60/135,870 25 May 1999 (25.05.1999) US 60/154,582 17 September 1999 (17.09.1999) US 60/174,256 03 January 2000 (03.01.2000) US <b>(60) Parent Application or Grant</b> THE PENN STATE RESEARCH FOUNDATION [/]; (). BENKOVIC, Stephen, J. [/]; (). SHAPIRO, Lucille [/]; (). BAKER, Stephen, J. [/]; (). WAHNON, Daphne, C. [/]; (). WALL, Mark [/]; (). BENKOVIC, Stephen, J. [/]; (). SHAPIRO, Lucille [/]; (). BAKER, Stephen, J. [/]; (). WAHNON, Daphne, C. [/]; (). WALL, Mark [/]; (). MONAHAN, Thomas, J. ; ().		<b>Published</b>
<b>(54) Title: DNA METHYLTRANSFERASE INHIBITORS</b> <b>(54) Titre: INHIBITEURS DE METHYLTRANSFERASE D'ADN</b>  <b>(57) Abstract</b> This invention provides broad-spectrum antibiotics that are inhibitors of bacterial adenine DNA methyltransferases.  <b>(57) Abrégé</b> L'invention concerne des antibiotiques à large spectre qui sont des inhibiteurs de méthyltransférases d'ADN d'adénine bactériens.		